



DESIGN STANDARDS FOR: RESTORATION SEEDING

1. Unless stated otherwise, seeding shall be composed of 1/3 sheep fescue, 1/3 chewings fescue, and 1/3 hard fescue. Percentages are by weight. The application rate shall be 125 lbs./ac, except on steep slopes where it shall be 150 lbs./ac.
2. Fertilizer shall be 20N-20P-20K with 50% nitrogen as sulfur-coated area. The application rate shall be 650 lbs./ac.
3. Tackifier shall be composed of wood fiber mulch and applied at a rate of 2000 lbs./ac. On steep slopes tackifier shall be EcoAegis bonded fiber matrix from Fiber Marketing in Renton, Washington, or an approved equivalent.
4. With prior approval by Parks, the following native seed mixes may be used:

Streambank Area Seed Mix

Scientific Name	Common Name	PLS lbs/ac
Beckmannia syzigachne	American sloughgrass	1.5
Bromus carinatus	California brome	4.0
Elymus glaucus	Blue wildrye	4.5
Glyceria occidentalis	Western mannagrass	3.0
Hordeum brachyantherus	Meadow barley	2.0
		15.0

Disturbed Area Seed Mix

Scientific Name	Common Name	PLS lbs/ac
Bromus carinatus	California brome	3.0
Elymus glaucus	Blue wildrye	4.0
Poa compressa	Canada wildrye	1.0
		8.0

Emergent Area Seed Mix

Scientific Name	Common Name	PLS lbs/ac
Beckmannia syzigachne	American sloughgrass	3.0
Elymus glaucus	Blue wildrye	6.0
Glyceria occidentalis	Western mannagrass	2.0
Deschampsia caespitosa	Tufted hairgrass	1.0
		12.0

END OF SECTION